



1. The function,  $y = 2 - 4 \cos(2x)$ , has a period of  $x = \pi$ . The function oscillates about  $y = 2$  with an amplitude of 4. It begins at a minimum at  $(0, -2)$ , goes to a maximum at  $(\pi/2, 6)$ , then ends its cycle at  $(\pi, -2)$ . The maxima occur at  $x = \pi/2, 3\pi/2$ . The graph of the function is below.



Problem 1



Problem 2

2. The function,  $y = 5 \sin(3x) - 4$ , has a period of  $x = 2\pi/3$ . The function oscillates about  $y = -4$  with an amplitude of 5. It begins at  $(0, -4)$ , goes to a maximum at  $(\pi/6, 1)$ , continues through  $(\pi/3, -4)$ , then reaches a minimum at  $(\pi/2, -9)$ , and ends its cycle at  $(2\pi/3, -4)$ . The maxima occur at  $x = \pi/6, 5\pi/6, 3\pi/2$ . The graph of the function is above.


3. The function,  $y(t) = 7 - 4 \cos(\frac{\pi}{8}(t - 5))$ , has a period of  $T = 16$ . The function oscillates about  $y = 7$  (vertical shift) with an amplitude of 4. The phase shift is  $\phi = 5$ . There is an absolute maximum at  $(t_{max}, y(t_{max})) = (13, 11)$ . There is an absolute minimum at  $(t_{min}, y(t_{min})) = (5, 3)$ . The graph of the function is below.

4. a. For the Logistic growth model,  $P_{n+1} = F(P_n) = 2.8P_n - 0.0005P_n^2$  with  $P_0 = 1000$ , then

$$P_1 = 2.8(1000) - 0.0005(1000)^2 = 2800 - 500 = 2300$$


$$P_2 = 2.8(2300) - 0.0005(2300)^2 = 3795$$

b. At equilibrium,  $P_e = P_n = P_{n+1}$ , so  $P_e = 2.8P_e - 0.0005P_e^2$  or  $P_e(1.8 - 0.0005P_e) = 0$ . One solution is  $P_e = 0$ , and the other equilibrium satisfies  $1.8 - 0.0005P_e = 0$  or  $P_e = \frac{1.8}{0.0005} = 3600$ . The derivative of the updating function is  $F'(P) = 2.8 - 0.001P$ . At  $P_e = 0$ ,  $F'(0) = 2.8 > 1$ , so this equilibrium is unstable with solutions monotonically growing away from  $P_e = 0$ . At  $P_e = 3600$ ,  $F'(3600) = 2.8 - 3.6 = -0.8 > -1$ , so the higher equilibrium is stable with solutions oscillating, but approaching  $P_e = 3600$ .



Problem 3

c. We see that  $F(P) = P(2.8 - 0.0005P)$ , so the updating function has  $P$ -intercepts at  $P = 0$  and  $P = 5600$ . The vertex has  $P_v = 2800$ , so  $F(2800) = 3920$ , which gives the vertex  $(2800, 3920)$ . The updating function intersects the identity function at the equilibria,  $(0, 0)$  and  $(3600, 3600)$ . The graph is shown below.



5. a. For the population model with the Allee effect,  $N_{n+1} = N_n + 0.1N_n \left(1 - \frac{1}{9}(N_n - 5)^2\right)$  with (population in thousands)  $N_0 = 4$ , the next two generations are

$$N_1 = 4 + 0.1(4) \left(1 - \frac{1}{9}(4 - 5)^2\right) = 4.356$$

$$N_2 = 4.356 + 0.1(4.356) \left(1 - \frac{1}{9}(4.356 - 5)^2\right) = 4.771$$

in thousands of birds.

b.  $N_e = N_e + 0.1N_e \left(1 - \frac{1}{9}(N_e - 5)^2\right)$ , so  $0.1N_e \left(1 - \frac{1}{9}(N_e - 5)^2\right) = 0$ . Thus,  $N_e = 0$  or  $(N_e - 5)^2 = 9$ . It follows that the equilibria are  $N_e = 0, 2$ , and  $8$ .

c. From the expanded model,  $N_{n+1} = A(N_n) = \frac{37}{45}N_n + \frac{1}{9}N_n^2 - \frac{1}{90}N_n^3$ , the derivative is  $A'(N) = \frac{37}{45} + \frac{2}{9}N - \frac{1}{30}N^2$ . At  $N_e = 0$ ,  $A'(0) = \frac{37}{45}$ , so this equilibrium is a stable equilibrium with solutions monotonically approaching 0. At  $N_e = 2$ ,  $A'(2) = \frac{17}{15}$ , so this equilibrium is an unstable equilibrium with solutions monotonically moving away from 2. At  $N_e = 8$ ,  $A'(8) = \frac{7}{15}$ , so this equilibrium is a stable equilibrium with solutions monotonically approaching 8.

d. Biologically, these results imply that if the population is below 2 thousand, then it will go to extinction ( $N_e = 0$ ). If the population is above 2 thousand, then the population of birds will grow to a carrying capacity of  $N_e = 8$  thousand.

6. The area of the brochure is  $A = xy = 125$ , where  $x$  is the width of the page and  $y$  is the length of the page. The area of the printed page, which is to be maximized is given by

$$P = (x - 4)(y - 5).$$

From the constraint on the page area, we have  $y = 125/x$ , which when substituted above gives

$$P(x) = (x - 4) \left( \frac{125}{x} - 5 \right) = 125 - \frac{500}{x} - 5x + 20 = 145 - 500x^{-1} - 5x.$$

The maximum is found by differentiation, which gives

$$P'(x) = 500x^{-2} - 5 = \frac{5(100 - x^2)}{x^2}.$$

This is zero when  $x = 10$ . It follows that  $y = 12.5$ . So the brochure has the dimensions  $10 \times 12.5$  with the printed region having dimensions  $6 \times 7.5$  or  $45 \text{ in}^2$ .

7. The volume of the open box satisfies the **Objective function**

$$V(x, y) = x^2y.$$

The **Constraint condition** on the surface area of this box is given by

$$SA = x^2 + 4xy = 600.$$

This constraint condition yields  $y = \frac{600 - x^2}{4x}$ , which when substituted into the objective function produces a function of one variable:

$$V(x) = x^2 \left( \frac{600 - x^2}{4x} \right) = \frac{1}{4}(600x - x^3).$$

Differentiating this quantity, we obtain

$$\frac{dV}{dx} = \frac{1}{4}(600 - 3x^2),$$

which when set equal to zero gives  $x = 10\sqrt{2}$ . (Take only the positive root.) This value of  $x$  gives the optimal length of one side of the base, which when substituted into the formula above gives  $y = 5\sqrt{2}$ . It follows that the maximum volume for this box is  $V(x) = 1000\sqrt{2}$ .

8. Combining the number of drops with the energy function, we have

$$E(h) = hN(h) = h \left( 1 + \frac{10}{h-1} \right) = h \left( \frac{h-1+10}{h-1} \right) = \frac{h^2 + 9h}{h-1}.$$

This is differentiated to give

$$E'(h) = \frac{(h-1)(2h+9) - (h^2+9h)}{(h-1)^2} = \frac{h^2 - 2h - 9}{(h-1)^2}.$$

A minimum occurs when  $h^2 - 2h - 9 = 0$ , so


$$h = 1 \pm \sqrt{10} = -2.1623, 4.1623.$$

It follows that the minimum energy occurs when  $h = 1 + \sqrt{10} = 4.1623$  m, which give the height that a crow should fly to minimize the energy needed to break open a walnut.


9. a. If  $P_0 = 100$ , then  $P_1 = 600e^{-0.1} = 542.9$  and  $P_2 = 6(542.9)e^{-0.5429} = 1892.75$ .

b. The derivative is  $R'(P) = 6e^{-0.001P}(1 - 0.001P)$ . The critical  $P_c$  occurs at  $P_c = 1000$ , so there is a maximum at  $(1000, 6000e^{-1}) = (1000, 2207)$ . The graph passes through the origin, so  $(0, 0)$  is the only intercept. Since  $\lim_{P \rightarrow \infty} R(P) = 0$ , there is a horizontal asymptote at  $R = 0$ . The second derivative is  $R''(P) = -0.006e^{-0.001P}(2 - 0.001P)$ , which is zero at  $P = 2000$ . Thus, there is a point of inflection at  $(2000, 12000e^{-2}) = (2000, 1624)$ . The graph is below.

c. The equilibria satisfy  $P_e = 6P_e e^{-0.001P_e}$ , so either  $P_e = 0$  or  $1 = 6e^{-0.001P_e}$ . The latter gives  $P_e = 1000 \ln(6) \simeq 1792$ . For  $P_e = 0$ ,  $R'(0) = 6 > 1$ , so this equilibrium is unstable with solutions moving monotonically away. For  $P_e = 1792$ ,  $R'(1792) = -0.7918$ , so this equilibrium is stable with solutions oscillating and moving toward the equilibrium.



Problem 9



Problem 10

10. a. If  $P_0 = 500$ , then  $P_1 = 8000/3.5^2 = 653.06$  and  $P_2 = 574.346$ .

b. The derivative is  $H'(P) = \frac{16(1 + 0.005P)^2 - 32P(1 + 0.005P)(0.005)}{(1 + 0.005P)^4} = \frac{16(1 - 0.005P)}{(1 + 0.005P)^3}$ .

The critical  $P_c$  occurs at  $P_c = 200$ , so there is a maximum at  $(200, 800)$ . The graph passes through the origin, so  $(0, 0)$  is the only intercept. Since  $\lim_{P \rightarrow \infty} H(P) = 0$ , there is a horizontal asymptote at  $H = 0$ . The second derivative is


$$H''(P) = \frac{-0.08(1 + 0.005P)^3 - 0.24(1 - 0.005P)(1 + 0.005P)^2}{(1 + 0.005P)^6} = \frac{-0.16(2 - 0.005P)}{(1 + 0.005P)^4},$$

which is zero at  $P = 400$ . Thus, there is a point of inflection at  $(400, 6400/9) = (400, 711)$ . The graph is above.

c. The equilibria satisfy  $P_e = 16P_e/(1 + 0.005P_e)^2$ , so either  $P_e = 0$  or  $(1 + 0.005P_e)^2 = 16$ . The latter gives  $P_e = 600$  (neglecting the negative solution). For  $P_e = 0$ ,  $H'(0) = 16 > 1$ , so this equilibrium is unstable with solutions moving monotonically away. For  $P_e = 600$ ,  $H'(600) = -0.5$ , so this equilibrium is stable with solutions oscillating and moving toward the equilibrium.

11. a. At rest,  $V(t) = -70 = 50t(t - 2)(t - 3) - 70$ , so  $50t(t - 2)(t - 3) = 0$ . Thus, the membrane is at rest when  $t = 0, 2$ , and  $3$ .

b. To find the extrema, we first write  $V(t) = 50(t^3 - 5t^2 + 6t) - 70$ , then the derivative is  $V'(t) = 50(3t^2 - 10t + 6)$ . By the quadratic formula,  $t = \frac{5}{3} \pm \frac{\sqrt{7}}{3} = 0.7847, 2.5486$ . Substituting these values into the membrane equation gives the peak of the action potential at  $t = 0.7847$  with a membrane potential of  $V(0.7847) = 35.63$  mV, while the minimum potential (most hyperpolarized state) occurs at  $t = 2.5486$  with a membrane potential of  $V(2.5486) = -101.56$  mV. Below is a graph for this model of membrane potential.



12. a. The time as a function of  $x$  is given by

$$T(x) = \frac{50 - x}{15} + \frac{(x^2 + 1600)^{1/2}}{9}.$$

b. We differentiate  $T(x)$  to find the minimum time,

$$T'(x) = -\frac{1}{15} + \frac{1}{9} \left( \frac{1}{2}(x^2 + 1600)^{-1/2} 2x \right) = -\frac{1}{15} + \frac{x}{9(x^2 + 1600)^{1/2}}.$$

Setting this derivative equal to zero gives

$$\begin{aligned} \frac{x}{9(x^2 + 1600)^{1/2}} &= \frac{1}{15} \\ 5x &= 3(x^2 + 1600)^{1/2} \\ 25x^2 &= 9(x^2 + 1600) \\ 16x^2 &= 14400 \\ x^2 &= 900 \end{aligned}$$

This implies  $x = 30$  m produces the minimum time.  $T(30) = \frac{20}{15} + \frac{50}{9} = \frac{62}{9} = 6.89$  sec. We check the endpoints  $T(0) = \frac{70}{9} = 7.778$  sec and  $T(50) = \frac{10\sqrt{41}}{9} = 7.11$  sec, confirming the optimal escape strategy is for the rabbit to run 20 m along the road, then run straight toward the burrow.

13. The **objective function** is given by:

$$S(x, y) = 2x^2 + 7xy.$$

The constraint condition is given by:

$$V = x^2y = 50,000 \text{ cm}^3, \quad \text{so,} \quad y = \frac{50,000}{x^2}.$$

Thus,

$$S(x) = 2x^2 + \frac{350,000}{x}.$$

Differentiating we have,

$$S'(x) = 4x - \frac{350,000}{x^2}.$$

Solving  $S'(x) = 0$ , so  $x^3 = \frac{350,000}{4} = 87,500$  or  $x = 44.395$ . It follows  $y = 25.37$ . Thus, the minimum amount of material needed is  $S(44.395) = 11,825.6 \text{ cm}^2$ .

14. From the diagrams, we have that  $r^2 + h^2 = a^2$ , which gives  $h^2 = a^2 - r^2$ . The circumference of the base of the cone is  $2\pi r = a\theta$ , where  $\theta$  is in radians. (Radians are an easy means of determining the length of a sector of a circle.) Thus,  $r = a\theta/2\pi$ . It follows that  $h^2 = a^2 - a^2\theta^2/(4\pi^2)$ . The volume of the water cup is given by

$$\begin{aligned} V &= \frac{\pi r^2 h}{3} = \frac{\pi}{3} \left( \frac{a\theta}{2\pi} \right)^2 \sqrt{a^2 - \frac{a^2\theta^2}{4\pi^2}} \\ V(\theta) &= \frac{a^3\theta^2}{12\pi} \sqrt{1 - \frac{\theta^2}{4\pi^2}} = \frac{a^3}{12\pi} \theta^2 \left( 1 - \frac{\theta^2}{4\pi^2} \right)^{1/2}. \end{aligned}$$

This expression is differentiated with respect to  $\theta$ .

$$V'(\theta) = \frac{a^3}{12\pi} \left( \frac{\theta^2}{2} \left( 1 - \frac{\theta^2}{4\pi^2} \right)^{-1/2} \left( \frac{-2\theta}{4\pi^2} \right) + 2\theta \left( 1 - \frac{\theta^2}{4\pi^2} \right)^{1/2} \right)$$

$$\begin{aligned}
&= \frac{a^3}{12\pi \left(1 - \frac{\theta^2}{4\pi^2}\right)^{1/2}} \left( \frac{-\theta^3}{4\pi^2} + 2\theta \left(1 - \frac{\theta^2}{4\pi^2}\right) \right) \\
&= \frac{a^3\theta}{12\pi \left(1 - \frac{\theta^2}{4\pi^2}\right)^{1/2}} \left( 2 - \frac{3\theta^2}{4\pi^2} \right)
\end{aligned}$$

The maximum is found by setting this derivative above equal to zero, so  $2 - \frac{3\theta^2}{4\pi^2} = 0$ . It follows that  $\theta^2 = 8\pi^2/3$  or

$$\theta = 2\pi\sqrt{\frac{2}{3}} \simeq 5.1302.$$

Thus,  $\theta = 2\pi\sqrt{\frac{2}{3}} \simeq 5.1302$  radians (which is about  $294^\circ$ ), so a sector of 1.1530 radians or about  $66^\circ$  is removed. The dimensions of the cone should have a radius of  $r = a\sqrt{\frac{2}{3}} \simeq 0.8165a$  and a height of  $h = a\sqrt{\frac{1}{3}} \simeq 0.57735a$ .

15. a.  $L(0) = 0.24$  m (24 cm) is the birth size a leopard shark ( $L$ -intercept). For large  $t$ ,  $L(t) \rightarrow 1.6$  m. The graph of this von Bertalanffy equation is shown below. Sexual maturity is found by solving  $L(t) = 0.5 = 1.6(1 - 0.85e^{-0.08t})$  or  $1.36e^{-0.08t} = 1.1$  or  $e^{0.08t} = 1.236$ . It follows that sexual maturity occurs at  $t = 2.652$  yr.

b. The composite function is given by

$$W(t) = 4.5(1.6(1 - 0.85e^{-0.08t}))^3 = 18.432(1 - 0.85e^{-0.08t})^3.$$

The intercept is  $W(0) = 0.0622$  kg, while for large  $t$ ,  $W(t) \rightarrow 18.432$  kg. The graph of this function is shown below.

c. By the chain rule, the derivative of  $W(t)$  is

$$W'(t) = 3(18.432)(1 - 0.85e^{-0.08t})^2(-0.85)(-0.08)e^{-0.08t} = 3.76e^{-0.08t}(1 - 0.85e^{-0.08t})^2.$$

By the product rule and chain rule, the second derivative is


$$\begin{aligned}
W''(t) &= 3.76 \left( 2e^{-0.08t}(1 - 0.85e^{-0.08t})(-0.85)(-0.08)e^{-0.08t} - 0.08e^{-0.08t}(1 - 0.85e^{-0.08t})^2 \right) \\
W''(t) &= 3.76e^{-0.08t}(1 - 0.85e^{-0.08t})(0.204e^{-0.08t} - 0.08)
\end{aligned}$$

$W''(t) = 0$  when either  $1 - 0.85e^{-0.08t} = 0$  or  $0.204e^{-0.08t} - 0.08 = 0$ . The first is zero when  $t = -2.03$  yr, while the second is zero when  $t = 11.7$  yr. It follows that the maximum weight gain occurs at age  $t = 11.7$  yr with a weight gain of  $W'(11.7) = 0.655$  kg/yr.


16. a. The periodic contractions of 10/min implies that  $0.1\omega = 2\pi$  or  $\omega = 20\pi$ . The average value  $A = \frac{4+1}{2} = 2.5$ , while the amplitude is given by  $B = 4 - 2.5 = 1.5$ . Thus, the radius of the small intestine is given by

$$R(t) = 2.5 + 1.5 \cos(20\pi t).$$

b. The graph of  $R(t)$  for  $t \in [0, 0.2]$  is shown below. The maxima occur at  $t = 0, 0.1, 0.2$  min, and the minima are halfway between the maxima with  $t = 0.05, 0.15$  min.



Problem 13a



Problem 13b

17. The period is 365 days, so  $365\omega = 2\pi$  or  $\omega = \frac{2\pi}{365} \simeq 0.01721$ . The average length of time is  $\alpha = \frac{1162+327}{2} = 744.5$  min. The amplitude is given by  $\beta = 1162 - 744.5 = 417.5$  min. The maximum occurs on day 170, so  $\omega(170 - \phi) = \pi/2$  (based on the maximum of the sine function). Thus,  $170 - \phi = \frac{365}{4} = 91.25$  or  $\phi = 78.75$  day. It follows that

$$L(t) = 744.5 + 417.5 \sin(0.01721(t - 78.75)).$$

The length of day for Ground Hog's day is  $L(32) = 744.5 + 417.5 \sin(0.01721(32 - 78.75)) = 443.7$  min in Anchorage.

18. a. From  $P_3$ , we have  $P_3 = 68.34 = 28.49(1+r)^3$ , so  $(1+r) = (68.34/28.49)^{1/3} = 1.33863$ . Thus,  $r = 0.33863$ . Doubling time satisfies  $2P_0 = P_0(1+r)^n$  or  $n = \ln(2)/\ln(1+r) = 2.377$  decades or 23.77 years.

b. The model predicts the population in 2000 is  $P_5 = 28.49(1.33863)^5 = 122.46$  million. The percent error is  $100 \frac{(122.46-99.93)}{99.93} = 22.55\%$ .


c. From the logistic model, we obtain  $P_1 = 39.32$  million and  $P_2 = 52.79$  million.

d. To find equilibria, we solve  $P_e = 1.48P_e - 0.0035P_e^2$ , which gives  $P_e = 0$  or  $P_e = 137.14$  million. The derivative of the updating function is  $F'(P) = 1.48 - 0.007P$ , so  $F'(137.14) = 0.52$ . It follows that this equilibrium is stable with solutions monotonically approaching this carrying capacity equilibrium.

19. From the high and low temperatures,  $A$  is the average, so  $A = 18^\circ\text{C}$ . The amplitude  $B$  is the difference between the maximum and the average, so  $B = 8^\circ\text{C}$ . The period is 24 hr, so  $24\omega = 2\pi$  or  $\omega = \frac{\pi}{12} \simeq 0.2618$ . The minimum temperature occurs at 4 AM ( $t = 4$ ), so

$$T(4) = 10 = 18 - 8 \sin\left(\frac{\pi}{12}(4 - \phi)\right).$$





It follows that

$$\sin\left(\frac{\pi}{12}(4 - \phi)\right) = 1 \quad \text{or} \quad \frac{\pi}{12}(4 - \phi) = \frac{\pi}{2}.$$

Hence,  $\phi = -2$ . If we want  $\phi \in [0, 24]$ , then by periodicity we can simply add 24 to obtain  $\phi = 22$ . (Both answers for  $\phi$  are correct, but if the restriction on  $\phi$  is required, we can only obtain the second answer.)

