

Math 531 - Partial Differential Equations

Fourier Series

Joseph M. Mahaffy,
 <jmahaffy@mail.sdsu.edu>

Department of Mathematics and Statistics
 Dynamical Systems Group
 Computational Sciences Research Center
 San Diego State University
 San Diego, CA 92182-7720

<http://jmahaffy.sdsu.edu>

Spring 2023



Introduction

The **separation of variables** technique solved our various **PDEs** provided we could write:

$$f(x) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos\left(\frac{n\pi x}{L}\right) + b_n \sin\left(\frac{n\pi x}{L}\right) \right).$$

Questions:

- 1 Does the infinite series converge?
- 2 Does it converge to $f(x)$?
- 3 Is the resulting infinite series really a solution of the PDE (and its subsidiary conditions)?

Mathematically, these are all difficult problems, yet these solutions have worked well since the early 1800's.



Outline

- 1 **Introduction**
 - Definitions
 - Convergence Theorem
 - Example
- 2 **Fourier Sine and Cosine Series**
 - Gibbs Phenomenon
 - Continuous Fourier Series
- 3 **Differentiation of Fourier Series**
 - Differentiation of Fourier Series
 - Differentiation of Cosine Series
 - Differentiation of Sine Series
- 4 **Method of Eigenfunction Expansion**



Definitions

Begin by restricting the class of $f(x)$ that we'll consider.

Definition (Piecewise Smooth)

A function $f(x)$ is **piecewise smooth** on some interval if and only if $f(x)$ is continuous and $f'(x)$ is continuous on a finite collection of sections of the given interval.

The only discontinuities allowed are jump discontinuities.

Definition (Jump Discontinuity)



A function $f(x)$ has a **jump discontinuity** at a point $x = x_0$, if the limit from the right $[f(x_0^+)]$ and the limit from the left $[f(x_0^-)]$ both exist and are not equal.

Piecewise smooth allows only a finite number of **jump discontinuities** in the function, $f(x)$, and its derivative, $f'(x)$.



Piecewise Smooth

The graph on the left is **piecewise smooth** with the function being continuous, but having a **jump discontinuity** in the derivative at $x = 0$




The graph on the right is **not piecewise smooth**, as the derivative becomes unbounded in any neighborhood of $x = 0$



Periodic Extension

The **Fourier series of $f(x)$** on an interval $-L \leq x \leq L$ is periodic with **period $2L$** .

However, the function $f(x)$ itself doesn't need to be periodic.



The graph above gives the **Fourier series period 2 extension** of $f(x) = x$ (along with $f(x)$, not periodic).



Fourier Series

Definitions of Fourier coefficients and a Fourier series. We must distinguish between a function $f(x)$ and its Fourier series over the interval $-L \leq x \leq L$.

$$\text{Fourier series} = a_0 + \sum_{n=1}^{\infty} a_n \cos\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi x}{L}\right).$$

The infinite series may not converge, and if it converges, it may not converge to $f(x)$

If the series converges, the **Fourier coefficients** a_0 , a_n , and b_n use certain **orthogonality integrals**.



Fourier coefficients

Definition (Fourier coefficients)

The definition of the **Fourier coefficients** are:

$$\begin{aligned} a_0 &= \frac{1}{2L} \int_{-L}^L f(x) dx \\ a_n &= \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx \\ b_n &= \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx \end{aligned}$$

The coefficients must be defined, e.g., $\left| \int_{-L}^L f(x) dx \right| < \infty$ for a_0 to exist. (No Fourier series for $f(x) = 1/x^2$.)



Fourier convergence

We write the **Fourier series**

$$f(x) \sim a_0 + \sum_{n=1}^{\infty} a_n \cos\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi x}{L}\right).$$

Theorem (Fourier convergence)

If $f(x)$ is **piecewise smooth** on the interval $-L \leq x \leq L$, then the **Fourier series** of $f(x)$ converges to:

- 1 The periodic extension of $f(x)$, where the periodic extension is continuous
- 2 The average of the two limits, usually $\frac{1}{2} [f(x^+) + f(x^-)]$, where the periodic extension has a **jump discontinuity**

Proof: The **proof of this theorem** requires significant techniques from Mathematical analysis, which is beyond the scope of this course. **SDSU**

Example

1

Example: Consider the Heaviside function shifted by 1:

$$f(x) = H(x-1) = \begin{cases} 0, & x < 1, \\ 1, & x \geq 1. \end{cases}$$

Find the Fourier series with $L = 2$.

The Fourier constant coefficient is

$$a_0 = \frac{1}{4} \int_{-2}^2 f(x) dx = \frac{1}{4} \int_1^2 1 dx = \frac{1}{4}.$$

The cosine coefficients:

$$\begin{aligned} a_n &= \frac{1}{2} \int_{-2}^2 f(x) \cos\left(\frac{n\pi x}{2}\right) dx = \frac{1}{2} \int_1^2 \cos\left(\frac{n\pi x}{2}\right) dx \\ &= \frac{\sin(n\pi) - \sin(n\pi/2)}{n\pi} = -\frac{1}{n\pi} \sin\left(\frac{n\pi}{2}\right). \end{aligned}$$

SDSU


Example

2


The sine coefficients:

$$\begin{aligned} b_n &= \frac{1}{2} \int_{-2}^2 f(x) \sin\left(\frac{n\pi x}{2}\right) dx = \frac{1}{2} \int_1^2 \sin\left(\frac{n\pi x}{2}\right) dx \\ &= \frac{\cos(n\pi/2) - \cos(n\pi)}{n\pi} = \frac{1}{n\pi} \left(\cos\left(\frac{n\pi}{2}\right) - (-1)^n \right). \end{aligned}$$

The **function**, $f(x)$, and **truncated Fourier series**.



Fourier series, $n = 20$



Fourier series, $n = 200$

SDSU

Example

3

```
1 % Periodic Fourier series, -2 < x < 2
2 % Step function at x = 1
3
4 NptsX=2000;           % number of x pts
5 Nf=200;               % number of Fourier terms
6 x=linspace(-5,5,NptsX);
7
8 a0=1/4;
9 a=zeros(1,Nf);
10 b=zeros(1,Nf);
11 f=a0*ones(1,NptsX);
12
13 for n=1:Nf
14     a(n) = -sin(n*pi/2)/(n*pi); % Fourier cosine ...
15     b(n) = (cos(n*pi/2)-cos(n*pi))/(n*pi); % ...
16     % Fourier sine coefficients
```

SDSU

Example

4

```

16     fn=a(n)*cos((n*pi*x)/2) + ...
        b(n)*sin((n*pi*x)/2); % Fourier function(n)
17     f=f+fn;
18 end
19 set(gca,'FontSize',16);
20 plot(x,f,'b-','LineWidth',1.5);
21 hold on
22 plot([-5,1],[0,0],'r-','LineWidth',1.5);
23 plot([1,5],[1,1],'r-','LineWidth',1.5);
24 xlabel('$x$', 'FontSize',16, 'FontName',fontlabs, ...
        'interpreter','latex');
25 ylabel('$y$', 'FontSize',16, 'FontName',fontlabs, ...
        'interpreter','latex');
26 axis on; grid;
27
28 print -depsc eg200-gr.eps
    
```



Fourier Sine Series

If $f(x)$ is an **odd function**, then $a_0 = a_n = 0$ and only the sine series remains:

$$b_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx.$$

This series appeared for solutions of the **heat equation**, $0 < x < L$ with $u(0, t) = u(L, t) = 0$

The **Sine series** produces an **odd extension** of $f(x)$

$$f(x) \sim \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), \quad 0 < x < L,$$

$$B_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx.$$



Fourier Cosine Series

If $f(x)$ is an **even function**, then $b_n = 0$ and only the cosine series remains:

$$f(x) \sim A_0 + \sum_{n=1}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), \quad 0 < x < L,$$

where

$$A_0 = \frac{1}{L} \int_0^L f(x) dx \quad \text{and} \quad A_n = \frac{2}{L} \int_0^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx.$$

This series appeared for solutions of the **heat equation**, $0 < x < L$ with $u_x(0, t) = u_x(L, t) = 0$.



Gibbs Phenomenon

1

Let $f(x) = 100$, and consider the **odd extension** of this function, so $f(x)$ is defined by

$$f(x) = \begin{cases} 100, & 0 < x < L, \\ -100, & -L < x < 0. \end{cases}$$

and extend it periodically with period $2L$.

As an **odd function**, this has a **Fourier sine series**

$$f(x) \sim \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right),$$

with

$$B_n = \frac{2}{L} \int_0^L 100 \sin\left(\frac{n\pi x}{L}\right) dx = \begin{cases} \frac{400}{n\pi}, & n \text{ odd,} \\ 0, & n \text{ even.} \end{cases}$$




Gibbs Phenomenon

2

We examine the graph for $n = 1, 3, 5, 7$ of

$$f(x) \sim \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), \quad \text{with } B_n = \begin{cases} \frac{400}{n\pi}, & n \text{ odd,} \\ 0, & n \text{ even.} \end{cases}$$




SDSU

Gibbs Phenomenon


3

We examine the graphs for $n = 40$ (20 nonzero terms) and $n = 200$ (100 nonzero terms) for

$$f(x) \sim \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), \quad \text{with } B_n = \begin{cases} \frac{400}{n\pi}, & n \text{ odd,} \\ 0, & n \text{ even.} \end{cases}$$



$n = 40$



$n = 200$

SDSU

Gibbs Phenomenon

4

The **Fourier series** for the $2L$ -periodic, **odd extension** of $f(x) = 100$,

$$f(x) \sim \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), \quad \text{with } B_n = \begin{cases} \frac{400}{n\pi}, & n \text{ odd,} \\ 0, & n \text{ even.} \end{cases}$$

It is clear that the **Fourier series** converges to **0** at $x = 0$ as every term in the series is **0**.

Similarly, the **Fourier series** converges to **0** at any $x = nL$ for $n = 0, \pm 1, \pm 2, \dots$, as every term in the series is also **0**.

The **Fourier Convergence Theorem** claims that the series converges to **100** for each $0 < x < L$.

SDSU

Gibbs Phenomenon

5

The $2L$ -periodic, **odd extension** of $f(x) = 100$,

$$f(x) \sim \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right), \quad \text{with } B_n = \begin{cases} \frac{400}{n\pi}, & n \text{ odd,} \\ 0, & n \text{ even.} \end{cases}$$

by the **Fourier Convergence Theorem** converges to **100** for $0 < x < L$, which is hard to show for most values of x .

Consider $x = \frac{L}{2}$,

$$\sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi}{2}\right) = \frac{400}{\pi} \left(1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots\right)$$

Euler's formula gives $\frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots$, (which is a very inefficient way to compute π , as it is an alternating series that does not **converge absolutely**)

SDSU

Gibbs Phenomenon

6

Harder to show convergence for other values of $x \in (0, L)$.

Convergence easily visualized as worst near **jump discontinuity**


For any finite sum in the series near $x = 0$, the solution starts at 0, then shoots up beyond 100, the primary overshoot

Examine previous $f(x)$

Figure (close up) with $n = 1000$ (or 500 nonzero terms)

The overshoot is about 20%

The maximum occurs at (0.01, 117.898)



SDSU

Gibbs Phenomenon

7

This **overshoot** is an example of the **Gibbs phenomenon**

For large n , in general, there is an overshoot of approximately 9% of the jump discontinuity

Note the previous example had a jump of **200**, and we saw the maximum of **117.898**, which is 9% of the jump

The **Gibbs phenomenon** only occurs for a finite series at a **jump discontinuity**

SDSU

Continuous Fourier Series

Theorem (Fourier Series)

For a piecewise smooth $f(x)$, the **Fourier series** of $f(x)$ is continuous and converges to $f(x)$ for $x \in [-L, L]$ if and only if $f(x)$ is continuous and $f(-L) = f(L)$.

Theorem (Fourier Cosine Series)

For a piecewise smooth $f(x)$, the **Fourier cosine series** of $f(x)$ is continuous and converges to $f(x)$ for $x \in [0, L]$ if and only if $f(x)$ is continuous.

Theorem (Fourier Sine Series)

For a piecewise smooth $f(x)$, the **Fourier sine series** of $f(x)$ is continuous and converges to $f(x)$ for $x \in [0, L]$ if and only if $f(x)$ is continuous and both $f(0) = 0$ and $f(L) = 0$.

SDSU

Differentiation of Fourier Series

Previously, we solved

$$\text{PDE: } \frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2}, \quad \text{BC: } u(0, t) = 0, \\ u(L, t) = 0.$$

IC: $u(x, 0) = f(x)$,
 and obtained the solution

$$u(x, t) = \sum_{n=1}^{\infty} B_n e^{-\frac{kn^2\pi^2t}{L^2}} \sin\left(\frac{n\pi x}{L}\right).$$

The **Superposition principle** justified this solution for any **finite series**, but can it be extended to the **infinite series**?

If $f(x)$ is piecewise smooth, then the **Fourier Convergence Theorem** shows that the **Fourier series** converges to the **Initial Conditions**

SDSU

Differentiation of Fourier Series

Suppose we can differentiate the series term-by-term, then in t

$$\frac{\partial u}{\partial t} = - \sum_{n=1}^{\infty} \frac{kn^2\pi^2}{L^2} B_n e^{-\frac{kn^2\pi^2 t}{L^2}} \sin\left(\frac{n\pi x}{L}\right).$$

Taking two partials with respect to x gives

$$\frac{\partial^2 u}{\partial x^2} = - \sum_{n=1}^{\infty} \frac{n^2\pi^2}{L^2} B_n e^{-\frac{kn^2\pi^2 t}{L^2}} \sin\left(\frac{n\pi x}{L}\right).$$

It follows that our solution above satisfies the **heat equation**:

$$u_t = ku_{xx}.$$



Counterexample

1

Differentiation Counterexample: Consider the **Fourier sine series** for $f(x) = x$ with $x \in [0, L]$:

$$x \sim \sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi x}{L}\right).$$

The **Fourier coefficients** satisfy:

$$\begin{aligned} b_n &= \frac{2}{L} \int_0^L x \sin\left(\frac{n\pi x}{L}\right) dx \\ &= \frac{2L}{n^2\pi^2} \left(\sin\left(\frac{n\pi x}{L}\right) - \frac{n\pi x}{L} \cos\left(\frac{n\pi x}{L}\right) \right) \Big|_0^L \\ &= -\frac{2L}{n\pi} \cos(n\pi) = \frac{2L}{n\pi} (-1)^{n+1} \end{aligned}$$

Thus, we have

$$x \sim \sum_{n=1}^{\infty} \frac{2L}{n\pi} (-1)^{n+1} \sin\left(\frac{n\pi x}{L}\right), \quad x \in [0, L].$$



Counterexample

2

Differentiation Counterexample: Continuing with

$$x \sim \sum_{n=1}^{\infty} \frac{2L}{n\pi} (-1)^{n+1} \sin\left(\frac{n\pi x}{L}\right), \quad x \in [0, L],$$

we differentiate the series term-by-term and obtain:

$$2 \sum_{n=1}^{\infty} (-1)^{n+1} \cos\left(\frac{n\pi x}{L}\right).$$

However, the series above is clearly not the cosine series for $f'(x) = 1$ (the derivative of x)

This series fails to converge anywhere, since the n^{th} term doesn't approach zero!



Differentiation of Fourier Series

When is term-by-term differentiation justified?

Theorem (Term-by-Term Differentiation)

A **Fourier series** that is continuous can be differentiated term-by-term if $f'(x)$ is **piecewise smooth**.

Corollary

If $f(x)$ is **piecewise smooth**, then the **Fourier series** of a continuous function, $f(x)$ can be differentiated term-by-term if $f(-L) = f(L)$.



Differentiation of Fourier Cosine Series

From our earlier result, if $f(x)$ is continuous, then its Fourier cosine series is continuous, avoiding *jump discontinuities* where difficulties occur for term-by-term differentiation

Theorem (Cosine Series Term-by-Term Differentiation)

If $f'(x)$ is *piecewise smooth*, then a continuous *Fourier cosine series* of $f(x)$ can be differentiated term-by-term.

Corollary (Cosine Series Term-by-Term Differentiation)

If $f(x)$ is *piecewise smooth*, then the *Fourier cosine series* of a continuous function $f(x)$ can be differentiated term-by-term.

Cosine Series Term-by-Term Differentiation

Thus, if

$$f(x) = A_0 + \sum_{n=1}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right), \quad 0 \leq x \leq L,$$

where equality implies convergence for all $0 \leq x \leq L$, the theorem above implies that

$$f'(x) \sim - \sum_{n=1}^{\infty} \left(\frac{n\pi}{L}\right) A_n \sin\left(\frac{n\pi x}{L}\right).$$

This sine series converges to points of continuity of $f'(x)$ and to the average where the Fourier sine series of $f'(x)$ is discontinuous.

Cosine Example

1

Example: Consider $f(x) = x$ on $0 \leq x \leq L$. Create an even extension, then make this $2L$ -periodic as seen in the graph.

The function has a continuous, piecewise smooth Fourier cosine series.


By our theorem, this *Fourier series* converges

The *Fourier coefficients* are

$$A_0 = \frac{1}{L} \int_0^L x dx = \frac{x^2}{2L} \Big|_0^L = \frac{L}{2}$$

and

$$\begin{aligned} A_n &= \frac{2}{L} \int_0^L x \cos\left(\frac{n\pi x}{L}\right) dx = \left(\frac{2L}{n^2\pi^2} \cos\left(\frac{n\pi x}{L}\right) + \frac{2x}{n\pi} \sin\left(\frac{n\pi x}{L}\right) \right) \Big|_0^L \\ &= \frac{2L}{n^2\pi^2} ((-1)^n - 1) \end{aligned}$$



Cosine Example

2

Thus,

$$x = \frac{L}{2} - \frac{4L}{\pi^2} \sum_{n \text{ odd}} \frac{1}{n^2} \cos\left(\frac{n\pi x}{L}\right),$$

where the series converges pointwise to the graph on the previous slide.


Note: This series converges absolutely by comparison to the series for $\frac{1}{n^2}$

The derivative of $f(x)$ is piecewise constant, as seen in the graph (right).

Differentiating term-by-term gives

$$1 \sim \frac{4}{\pi} \sum_{n \text{ odd}} \frac{1}{n} \sin\left(\frac{n\pi x}{L}\right), \quad 0 < x < L.$$

The weaker series convergence is easily seen, and it is easy to verify that this is the sine series for $f'(x) = 1$.



Sine Series Term-by-Term Differentiation

Similar results hold for the **sine series** with more conditions

Theorem

Sine Series Term-by-Term Differentiation] If $f'(x)$ is **piecewise smooth**, then a continuous **Fourier sine series** of $f(x)$ can be differentiated term-by-term.

Corollary (Sine Series Term-by-Term Differentiation)

If $f'(x)$ is **piecewise smooth**, then the **Fourier sine series** of a continuous function $f(x)$ can be differentiated term-by-term if $f(0) = 0$ and $f(L) = 0$.



Sine Series Term-by-Term Differentiation

Proof: We prove term-by-term differentiation of the **Fourier sine series** of a **continuous** function $f(x)$, when $f'(x)$ is **piecewise smooth** and $f(0) = 0 = f(L)$:

$$f(x) \sim \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right),$$

where B_n are expressed later. Equality holds if $f(0) = 0 = f(L)$.

If $f'(x)$ is piecewise smooth, then $f'(x)$ has a **Fourier cosine series**

$$f'(x) \sim A_0 + \sum_{n=1}^{\infty} A_n \cos\left(\frac{n\pi x}{L}\right),$$

where A_0 and A_n are expressed later.

This series will not converge to $f'(x)$ at points of **discontinuity**.



Sine Series Term-by-Term Differentiation

Proof (cont): Need to verify that

$$f'(x) \sim \sum_{n=1}^{\infty} \frac{n\pi}{L} B_n \cos\left(\frac{n\pi x}{L}\right).$$

The **Fundamental Theorem of Calculus** gives:

$$A_0 = \frac{1}{L} \int_0^L f'(x) dx = \frac{1}{L} (f(L) - f(0)).$$

Integrating by parts,

$$\begin{aligned} A_n &= \frac{2}{L} \int_0^L f'(x) \cos\left(\frac{n\pi x}{L}\right) dx \\ &= \frac{2}{L} \left[f(x) \cos\left(\frac{n\pi x}{L}\right) \Big|_0^L + \frac{n\pi}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx \right] \end{aligned}$$



Sine Series Term-by-Term Differentiation

Proof (cont): However, B_n , the **Fourier sine series coefficient** of $f(x)$ is

$$B_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx,$$

so for $n \neq 0$

$$A_n = \frac{n\pi}{L} B_n + \frac{2}{L} \left[(-1)^n f(L) - f(0) \right].$$

It follows that we need $f(0) = 0 = f(L)$ for both $A_0 = 0$ and $A_n = \frac{n\pi}{L} B_n$, completing the proof.

However, this proof gives us more information about **differentiating the Fourier sine series**.



Sine Series Term-by-Term Differentiation

The more general theorem for *differentiating the Fourier sine series* is below:

Theorem

If $f'(x)$ is *piecewise smooth*, then the *Fourier sine series* of a continuous function $f(x)$,

$$f(x) \sim \sum_{n=1}^{\infty} B_n \sin\left(\frac{n\pi x}{L}\right)$$

cannot, in general be differentiated term-by-term. However,

$$f'(x) \sim \frac{1}{L} \left[f(L) - f(0) \right] + \sum_{n=1}^{\infty} \left(\frac{n\pi}{L} B_n + \frac{2}{L} \left[(-1)^n f(L) - f(0) \right] \right) \cos\left(\frac{n\pi x}{L}\right).$$



Sine Series Term-by-Term Differentiation

Example: Previously considered $f(x) = x$ with a *Fourier sine series* and showed this could not be differentiated term-by-term.

The *Fourier sine series* satisfies:

$$f(x) = x \sim 2 \sum_{n=1}^{\infty} \frac{L(-1)^{n+1}}{n\pi} \sin\left(\frac{n\pi x}{L}\right).$$

Since $f(0) = 0$ and $f(L) = L$, from the general formula above:

$$A_0 = \frac{1}{L} \left(f(L) - f(0) \right) = 1.$$

and

$$\begin{aligned} A_n &= \frac{n\pi}{L} B_n + \frac{2}{L} \left[(-1)^n f(L) - f(0) \right] \\ &= 2(-1)^{n+1} + 2(-1)^n = 0. \end{aligned}$$

It follows that we obtain the correct derivative

$$f'(x) = 1.$$



Method of Eigenfunction Expansion

Want to apply techniques of *differentiating a Fourier series* term-by-term to **PDEs**

Use an alternative **method of eigenfunction expansion**, which can be applied to **nonhomogeneous BCs**

Consider an *eigenfunction expansion* of the form

$$u(x, t) \sim \sum_{n=1}^{\infty} B_n(t) \sin\left(\frac{n\pi x}{L}\right),$$

where the *Fourier sine coefficients* depend on time, t



Method of Eigenfunction Expansion

The initial condition, $u(x, 0) = f(x)$, is satisfied if

$$f(x) \sim \sum_{n=1}^{\infty} B_n(0) \sin\left(\frac{n\pi x}{L}\right),$$

where the initial *Fourier sine coefficients* are

$$B_n(0) = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

Can we differentiate term-by-term to satisfy the **heat equation**,

$$u_t = k u_{xx}?$$

Need **two** partial derivatives with respect to x and **one** partial derivative with respect to t .



Method of Eigenfunction Expansion

If $u(x, t)$ is continuous, then the *Fourier sine series* can be differentiated term-by-term provided

$$u(0, t) = 0 \quad \text{and} \quad u(L, t) = 0.$$

(homogeneous BCs)

The result is

$$\frac{\partial u}{\partial x} \sim \sum_{n=1}^{\infty} \frac{n\pi}{L} B_n(t) \cos\left(\frac{n\pi x}{L}\right),$$

which is a *Fourier cosine series*

Provided $\frac{\partial u}{\partial x}$ is continuous, it can be differentiated term-by-term:

$$\frac{\partial^2 u}{\partial x^2} \sim - \sum_{n=1}^{\infty} \frac{n^2 \pi^2}{L^2} B_n(t) \sin\left(\frac{n\pi x}{L}\right),$$



Method of Eigenfunction Expansion

The **two** derivatives w.r.t. x could be taken term-by-term provided the problem has homogeneous BCs.

Need

$$\frac{\partial u}{\partial t} \sim \sum_{n=1}^{\infty} \frac{dB_n}{dt} \sin\left(\frac{n\pi x}{L}\right).$$

If term-by-term evaluation is justified, then

$$\frac{dB_n}{dt} = -k \frac{n^2 \pi^2}{L^2} B_n(t),$$

so

$$B_n(t) = B_n(0) e^{-\frac{n^2 \pi^2}{L^2} kt}.$$



Method of Eigenfunction Expansion

Theorem

The *Fourier series* of a continuous function $u(x, t)$

$$u(x, t) = a_0(t) + \sum_{n=1}^{\infty} \left(a_n(t) \cos\left(\frac{n\pi x}{L}\right) + b_n(t) \sin\left(\frac{n\pi x}{L}\right) \right),$$

can be differentiated term-by-term with respect to t

$$\frac{\partial u(x, t)}{\partial t} = a_0'(t) + \sum_{n=1}^{\infty} \left(a_n'(t) \cos\left(\frac{n\pi x}{L}\right) + b_n'(t) \sin\left(\frac{n\pi x}{L}\right) \right),$$

if $\frac{\partial u}{\partial t}$ is *piecewise smooth*.

This theorem justifies the use of separation of variables and our solution.



Term-by-Term Integration

Theorem

A *Fourier series* of a piecewise smooth $f(x)$ can always be integrated term-by-term and the result is a convergent infinite series that always converges to the integral of $f(x)$ for $-L \leq x \leq L$ (even if the original Fourier series has *jump discontinuities*).

