

Math 534A Solution to Problem 0.12

As promised at my office hour last week, here's the solution to problem 12 from the Introduction in Marsden & Hoffmann.

Problem: Let  $\mathcal{A}$  be a collection of subsets of a set  $S$  and  $\mathcal{B}$  the collection of complementary sets; that is,  $B \in \mathcal{B}$  iff  $S \setminus B \in \mathcal{A}$ . Prove de Morgan's laws:

a.  $S \setminus \bigcup \mathcal{A} = \bigcap \mathcal{B}$ .

Proof: We begin by noting that

$$\bigcup \mathcal{A} = \{a \in S \mid \exists A \in \mathcal{A}, a \in A\}. \quad (1)$$

Then by the rules of negation,

$$S \setminus (\bigcup \mathcal{A}) = \{a \in S \mid \forall A \in \mathcal{A}, a \notin A\} \quad (2)$$

But

$$\bigcap \mathcal{B} = \{a \in S \mid \forall B \in \mathcal{B}, a \in B\} \quad (3)$$

and by using the definition of  $\mathcal{B}$  this becomes

$$\bigcap \mathcal{B} = \{a \in S \mid \forall (S \setminus B) \in \mathcal{A}, a \in B\} \quad (4)$$

or

$$\bigcap \mathcal{B} = \{a \in S \mid \forall (S \setminus B) \in \mathcal{A}, a \notin (S \setminus B)\}. \quad (5)$$

Renaming  $S \setminus B$ ,  $A$  gives the result. This is justified since in the condition  $S \setminus B$  runs over all subsets of  $S$ .

b.  $S \setminus \bigcap \mathcal{A} = \bigcup \mathcal{B}$ .

Proof: Noting that  $S \setminus (S \setminus X) = X$  for all subsets  $X$  of  $S$ , this follows by taking complements in part a, and switching the roles of the two collections.