

Math 254 Exam 4 Solutions

1. Carefully state the definition of “subspace”. Give two examples, each from \mathbb{R}^2 .

A subspace is a subset of a vector space that is, itself, a vector space. Many examples are possible, such as $\{(0, 0)\}$ (zero-dimensional), $\text{Span}(S)$ for $S = \{(1, 2)\}$ (one-dimensional), or \mathbb{R}^2 itself (two-dimensional).

2. Carefully state any five of the eight vector space axioms.

These are listed on p.152 of the text. It is not important how you number them; however it is important that you give the English text correctly. “For any vectors u, v, w in V , $(u + v) + w = u + (v + w)$.” is correct, but the equation “ $(u + v) + w = u + (v + w)$ ” alone is incorrect.

3. Let $S = \{f(x) : f(3) = 1\} \subseteq \mathbb{R}[x]$ be the set of all polynomials $f(x)$ satisfying $f(3) = 1$. Determine, with justification, whether this is a vector space.

Since S is a subset of a vector space, to be a subspace S must satisfy three properties. It must contain the zero vector, it must be closed under vector addition, and closed under scalar multiplication. S satisfies *none* of these three properties, and it's enough to pick your favorite to disprove. Just for fun, I will disprove all three: (1) $f(x) = 0$ does not satisfy $f(3) = 1$, so 0 is not in S ; (2) $f(x) = 1$ and $g(x) = x/3$ are both in S , but $(f+g)(x) = 1 + x/3$ is not in S since $(f+g)(3) = 2$; (3) $f(x) = 1$ is in S but $5f(x) = 5$ is not in S .

4. Determine, with justification, whether $(1, 2)$ is in the row space of $M = \begin{bmatrix} 2 & 3 \\ 6 & 9 \end{bmatrix}$.

The row space of M is also the row space of $\begin{bmatrix} 2 & 3 \\ 0 & 0 \end{bmatrix}$, obtained via $R_2 = R_2 - 3R_1$, which is $\text{Span}((2, 3)) = \{t(2, 3) : t \in \mathbb{R}\}$. If $(1, 2)$ were in this subspace, then for some t we would have $(1, 2) = (2t, 3t)$, and hence $2t = 1$ and $3t = 2$. This is impossible, so the answer is “no”.

5. Set $V = \mathbb{R}^3$. Give any two subspaces U_1, U_2 such that $U_1 \oplus U_2 = V$.

Two type of solutions are possible. The “trivial” solution is $U_1 = \{(0, 0, 0)\}$, $U_2 = \mathbb{R}^3$ (or the other way around). Otherwise, one of U_1, U_2 will be one-dimensional and the other will be two-dimensional. Many examples are possible, for example $U_1 = \text{Span}(\{(1, 0, 0)\}) = \{(a, 0, 0) : a \in \mathbb{R}\}$, $U_2 = \text{Span}(\{(0, 1, 0), (0, 0, 1)\}) = \{(0, b, c) : b, c \in \mathbb{R}\}$. What is important is that U_1, U_2 are both subspaces of \mathbb{R}^3 , that $U_1 + U_2 = \mathbb{R}^3$, and that $U_1 \cap U_2 = \{0\}$.